

Impacts of a reduction of seawater pH mimicking ocean acidification impacts on assemblage, structure and diversity of marine fungal communities

Reich, Marlis; Wichels, Antje; Panzer, Katrin ; Krause, Evamaria; Gimenez Noya, Jose; Gerdts, Gunnar

Aquatic Microbial Ecology

DOI:
[10.3354/ame01831](https://doi.org/10.3354/ame01831)

Published: 01/01/2017

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Reich, M., Wichels, A., Panzer, K., Krause, E., Gimenez Noya, J., & Gerdts, G. (2017). Impacts of a reduction of seawater pH mimicking ocean acidification impacts on assemblage, structure and diversity of marine fungal communities. *Aquatic Microbial Ecology*, 79(3), 221-233. <https://doi.org/10.3354/ame01831>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Impacts of a reduction in seawater pH mimicking ocean acidification on the structure and diversity of mycoplankton communities

Marlis Reich^{1*}, Antje Wichels², Katrin Panzer^{1,3}, Evamaria Krause⁴, Luis Giménez⁵, Gunnar Gerds².

¹Molecular Ecology, Institute of Ecology, FB02, University of Bremen, Bremen, Germany

²Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Biologische Anstalt Helgoland, PO Box 180, 27498 Helgoland, Germany

³Jacobs University Bremen gGmbH, Department of Life Sciences and Chemistry, Campusring 1, 28759 Bremen, Germany

⁴Department of Natural Sciences, Mathematics and Medicine, BIS – Library and Information Systems, University of Oldenburg, Oldenburg, Germany

⁵School of Ocean Sciences, Bangor University UK, Askew Street, Menai Bridge, Anglesey LL59 5AB, UK

Corresponding author:

Marlis Reich, University of Bremen, Molecular Ecology, FB02, Leobener Str. 2, Bremen, 28359, Germany, Phone: +49-(0)421-21862825, Fax: +49-(0)421-2189862825, Mail: reich@uni-bremen.de

Short title: Ocean acidification impacts fungal community structure and diversity

Abstract

Increases in atmospheric carbon dioxide (CO₂) change ocean chemistry, as dissolved CO₂ leads to a reduction in the seawater pH. Many marine taxa have been shown to be affected by ocean acidification, while information on marine fungi is lacking. Here, we analyze the effect of pH on mycoplankton communities. The pH of microcosms was adjusted to a value mimicking the predicted ocean acidification in the near future. Fungal communities were analyzed using a double-marker gene approach, allowing a more detailed analysis of their response using 454 pyrosequencing. Mycoplankton communities in microcosms with *in situ* and adjusted water pH values differed significantly in terms of structure and diversity. The differences were mainly based on abundance shifts among the dominant taxa rather than the exclusion of fungal groups. A sensitivity to lower pH values was reported for several groups across the fungal kingdom and was not phylogenetically conserved. Some of the fungal species that dominated the communities of microcosms with a lower pH were known pathogenic fungi. With the increasing awareness of the significant role fungi play in marine systems, including performing a diverse range of symbiotic activities, our results highlight the importance of including fungi in further research projects studying and modeling biotic responses to the predicted ocean acidification.

Keywords: ocean acidification, marine fungi, phylogenetic signals, 18S rRNA gene sequence, ITS, microcosm, double-marker gene approach

1. Introduction

Oceans cover approximately 72% of the Earth's surface and are home to an uncountable number of organisms. However, over the last two centuries, environmental conditions in ocean systems have been changing due to an enormous increase in atmospheric carbon dioxide (CO₂) caused by anthropogenic activities, such as the burning of fossil fuels (Sabine et al. 2004). The oceans are known to act as a net sink for atmospheric CO₂. Dissolution of CO₂ in water results in the formation of carboxylic acid and therefore seawater acidification (Caldeira & Wickett 2003). This mechanism, called ocean acidification, has led to a drop in seawater pH by 0.1 units since the industrial revolution (Sabine et al. 2004). Future scenarios predict a further decrease by 0.4 units within the current century (Caldeira & Wickett 2005).

Many organism groups seem to be sensitive to such a drop in pH, as reported for foraminifera (Webster et al. 2013), animals (Harvell et al. 2002, Hoegh-Guldberg et al. 2007), and bacteria (Witt et al. 2011, Krause et al. 2012). In contrast to other marine microorganisms, the importance of fungi for marine ecosystem functioning has long been underappreciated, which is why they have been highly understudied. Only in recent years, the use of next-generation sequencing (NGS) techniques has shed some light on fungal diversity (Richards et al. 2015, Rama et al. 2016, Taylor & Cunliffe 2016) and functioning (Gutierrez et al. 2011, Orsi et al. 2013, Orsi et al. 2015). Fungal diversity assessments are generally conducted using internal transcribed spacer (ITS) regions as marker genes (Schoch et al. 2012). However, the ITS is not the best marker for all fungal groups, and many environmental sequences can often only be classified to the phylum or kingdom level (Nilsson et al. 2016). To achieve a better resolution of fungal communities, Arfi *et al.* (2012) proposed to include a second phylogenetic marker (18S rDNA). High-resolution studies on marine fungal communities and their responses to ocean acidification are currently lacking, although fungi can be affected by changes in pH in diverse ways; one example is the uptake of nutrients and organic compounds driven by the

46 electrochemical gradient of H⁺ ions across the plasma membrane (Bowman & Bowman 1986). The
47 activity and kinetics of excreted fungal exoenzymes are also mediated by the medium pH (Pritsch et
48 al. 2004, Courty et al. 2005). A further but indirect effect is the availability and accessibility of
49 substrates, such as the range and size of dissolved and particulate organic matter (Verdugo et al.
50 2004).

51 In a previous study, we tested the pH sensitivity of marine fungal communities in microcosms using
52 ITS fingerprinting (automated ribosomal intergenic spacer analysis; ARISA) (Krause et al. 2013).
53 The seawater pH was altered according to one of the possible near-future scenarios (Blackford &
54 Gilbert 2007). Interestingly, the structure of the marine fungal communities showed a significant pH-
55 dependent response. However, details on the taxonomic composition of the communities and the taxa
56 involved in these structural changes could not be provided due to the inherent limitations of the
57 applied fingerprinting technique. Thus, the purpose of the present study was to investigate the
58 underlying dynamics of the fungal community composition. We tested the modified double-marker
59 gene approach of Arfi *et al.* (2012) by applying a phylogenetic-based analysis using 18S rDNA in
60 addition to a hierarchical cluster-based approach for the ITS. We aimed to identify which parameters
61 of the communities change, how they change, whether specific fungal species benefit from changes
62 in pH, and whether the observed sensitivity of fungi toward lower pH values is clade-specific. We
63 hypothesize that the observed shift in the community structure is based on the response of a few
64 fungal species and that pH sensitivity is phylogenetically conserved.

2. Materials and Methods

2.1. Establishment of the microcosm experiments and sampling scheme

Surface seawater samples were collected at the Helgoland Roads station in the North Sea (Germany) on the 3rd of May, 2012. The sampling site is located off the island of Helgoland in the southern North Sea (German Bight). Due to strong tidal currents, surface water samples are representative of the entire water body (Wiltshire et al. 2008). For the year 2100, a concentration of 1000 ppm of atmospheric CO₂ has been predicted (IPCC 2014), which would result in a mean surface seawater pH of 7.67 in the southern North Sea according to the model of Blackford & Gilbert (2007). Thus, after the sampled seawater was poured into 1.6 l glass jars, the microcosms were incubated either at the current *in situ* seawater pH (8.26) or at an adjusted pH of 7.67. The seawater pH was adjusted with 2 M HCl, which is a validated approach for acidifying microcosm-based experiments (Krause et al. 2012). For each pH treatment, 20 replicates were prepared and left in the dark over a maximum period of 4 weeks at the *in situ* temperature on the day of sampling (8°C). The microcosms were mixed daily by manual inversion. Every week, the biomass in 5 replicates for each pH treatment was collected by filtration onto sterile nitrocellulose filters (0.45 µm pore size, 47 mm diameter, gray with a grid; Sartorius, Göttingen, Germany). Parameters of the carbonate system of every microcosm, including the pH (Suppl. Table S1), were calculated based on a previously described method (Krause et al. 2013) and can be accessed at <https://doi.pangaea.de/10.1594/PANGAEA.831726> (Pangaea ID 10.3354/ame01622). Biomass filters were stored at -20°C until further treatment.

2.2. DNA extraction, PCR and pyrosequencing

DNA was extracted as described by Krause *et al.* (2013). The PCR on the ITS2 region was conducted in a reaction of 50 µl containing 100 ng of template DNA, 5 µl Taq buffer (10×), 2.5 µM MgCl₂, 250 µM of each dNTP, 2.5 U Taq polymerase (5 Prime, Germany), 0.15 µM of the forward fungal-specific primer gITS7 (5'GTGARTCATCGARTCTTTG) (Ihrmark et al. 2012) and 0.25 µM of the

89 tagged eukaryotic ITS4 primer (5'TCCTCCGCTTATTGATATGC) (White et al. 1990). Cycling
90 conditions followed Krause *et al.* (Krause et al. 2013). A second amplification round was conducted
91 in 2 µl of a 1:10 dilution of the first PCR products and using the primers described above. The 18S
92 rDNA was amplified in reactions of 50 µl containing 50 ng of the template DNA, 5 µl Taq buffer
93 (10×) (5 Prime), 1.25 µM MgCl₂, 200 µM of each dNTP, 0.3 µM each of the eukaryotic primer NS1
94 (5'GTAGTCATATGCTTGTCTC) and the fungal-specific primer NS8
95 (5'TCCGCAGGTTACCTACGGA) (White et al. 1990) and 1.25 U Taq polymerase. Cycling
96 conditions started with a denaturation step of 94°C for 15 min, followed by 30 cycles of 94°C for 1
97 min, 50°C for 1 min and 68°C for 2 min, with a final extension at 68°C for 10 min. For the second
98 round, the fungal-specific primer pair nu-SSU-0817 (5'TTAGCATGGAATAATRRATAGGA)/nu-
99 SSU-1536 (5'ATTGCAATGCYCTATCCCCA) (Borneman & Hartin 2000) was used, with the first
100 primer carrying the barcode. Amplifications were conducted in reactions of 50 µl, each containing 2
101 µl of the first PCR products, 5 µl Taq buffer (10×), 1.25 mM MgCl₂, 250 µM of each dNTP, 0.48
102 µM of each primer and 2.5 U Taq polymerase (5 Prime). Cycling conditions were the same as for the
103 first PCR but with an annealing temperature of 68°C. The PCR steps included a positive control,
104 which was a seawater sample taken at the time of sampling. After purification using the PeqGold Gel
105 Extraction kit (PeqLab, Erlangen, Germany), the PCR products were sequenced on a Roche 454 GS-
106 FLX Titanium platform (LGC, Berlin, Germany) as single reads. Sequencing data are stored under
107 the study accession number SRP065054 and can be downloaded from the Sequence Read Archive
108 (SRA) of the National Center for Biotechnology Information (NCBI) website
109 (<http://www.ncbi.nlm.nih.gov/sra>).

110 **2.3. Bioinformatics and OTU designation**

111 For quality control, ITS2 and 18S rDNA sequences without a valid primer or DNA tag sequence and
112 with average quality values of <25 Phred score, >2% ambiguous symbols or homopolymer chains of

113 >8 nt were discarded. After removing barcodes and primers from the remaining sequences, sequences
114 with a read length <200 nt were also discarded. In most sequence reads, quality values dropped
115 below a value of 25 at an nt position of ~ 420. Thus, all sequences were trimmed from the 3'-end to a
116 maximal length of 400 nt. Potential chimeric sequences were detected and removed using the
117 UCHIME program (Edgar et al. 2011). From this point on, the remaining ITS2 sequences were
118 subjected to hierarchical clustering analyses, while 18S rDNA sequences were further analyzed using
119 a phylogenetic-based approach.

120 **2.4. Fungal taxonomy**

121 Due to the primer attachment sites, the generated ITS reads contained sequence information from the
122 5.8S and 28S rDNA. As these fragments influence the sequence identification process, they were
123 removed from the generated sequence reads using the ITSx program (Bengtsson-Palme et al. 2013).
124 ITS2 sequences were clustered using a 97% sequence similarity threshold into operational taxonomic
125 units (OTUs) using the UCLUST algorithm (Edgar 2010). All these steps were processed using the
126 Quantitative Insights In Microbial Ecology (QIIME) pipeline (Caporaso et al. 2010). Sequences were
127 blasted using the FHiTINGS program (Dannemiller et al. 2014). Finally, OTUs that were represented
128 by fewer than five reads and observed only in a single sample were removed. 18S rDNA reads
129 passing the above-described quality control requirements were clustered and classified by mapping
130 the sequences to our existing maximum likelihood reference tree (for details on the sequences,
131 alignment and methods see Panzer *et al.* (2015). The effects of pH and time on specific fungal groups
132 were tested using generalized linear modeling (Zuur et al. 2009), as our data followed Poisson or
133 negative binomial distributions. Model selection was carried out using the Akaike information
134 criterion (AIC); models were ordered according to the AIC, and those with lowest scores were
135 compared using likelihood ratio tests (LR). The best model was then evaluated for significance of the
136 interaction term using LR. Tests were run in R (R Core Team 2015) using the “glm” and “glm.nb”

functions of the Mass library (Venables & Ripley 2002). The values from the “summary” table are reported.

2.5. Alpha diversity analyses based on the ITS2 dataset

Diversity indices were calculated based on subsampled datasets using the PRIMER 6 software (Plymouth Marine Laboratories, UK). Next, the effects of the factors “pH” and “incubation time” on the diversity indices were analyzed. In the case for the Simpson and Shannon diversity data, the statistical general least squares (GLS) model (Galecki & Burzykowski 2013) was chosen, which can deal with data with heterogeneous variances. The details of the statistical procedure are given in Zuur *et al.* (2009) and Galecki & Burzykowski (2013), and here we only describe them in brief. In the first stage, we fitted models with different variance structure (i.e., different patterns of variance heterogeneity) using restricted maximum likelihood (REML) and selected the model with the best variance structure using the AIC and LR as explained in section 2.4. In the second step, we used the model with the best variance structure to test for the effects of “pH” and “incubation time” on the average diversity. In this step, models are fitted using the maximum likelihood method (ML) and are then compared using the AIC, and significant effects are tested using LR. When significant effects are found, the parameters in the model with the significant terms are refitted using REML in order to express the appropriate estimates of variances and average trends. GLS analysis was carried out using the package “nlme” (Pinheiro *et al.* 2015) implemented in R (R Core Team 2015). The effects of pH and incubation time on OTU richness were evaluated using generalized linear modeling (Zuur *et al.* 2009) with Poisson and negative binomial error structure as in the case of species counts (see section 2.4).

2.6. Beta diversity analyses based on the ITS2 dataset

Beta diversity was analyzed based on subsampled datasets using permutational multivariate ANOVA (PERMANOVA) of square root-transformed OTU numbers (as Bray-Curtis similarities). To visualize the influence of the factors “pH” and “incubation time” on the fungal community structure, we performed principal coordinate analyses (PCoAs). The similarity percentage (SIMPER) procedure was used to identify OTUs that contributed most to the observed dissimilarity between the two pH treatments. We examined only OTUs with a threshold of > 1% of the total contribution to differences. All analyses were performed using the PRIMER 6 software.

2.7. Phylogenetic diversity of the 18S rDNA dataset

To evaluate the evolutionary relationships among taxa, we ran further phylogenetic metric-based PCoAs using the 18S rDNA dataset. Phylogenetic separation patterns were identified using the FastUniFrac suite implemented in the Galaxy platform (Giardine et al. 2005, Goecks et al. 2010). Unweighted and weighted distances were calculated.

2.8. Phylogenetic signals in the 18S rDNA dataset

To test for possible genetic conservatism among fungi to acclimate to the factor “pH”, phylogenetic signal analysis was performed using the software PHYLOCOM v 4.2 (Webb et al. 2008) with the Comstruct function. Evolutionary conservatism was analyzed based on the net relatedness index (NRI) and the nearest taxon index (NTI) using the 18S rDNA dataset. The NRI indicates how many taxa in a community are dispersed (negative value) or clustered (positive value) over the whole phylogenetic tree, while the NTI indicates dispersal/clustering at lower taxonomic levels. As a null model, the “-m 2” function was chosen, while the number of randomizations was set to 9999. A p-value of $p < 0.05$ against the random distributions was considered to indicate a significant phylogenetic signal. In a last step, Nodessig analyses were run to identify the clades/taxa in the phylogenetic tree that caused the phylogenetic signals.

3. Results

3.1. Dataset description

In total, 237,967 and 532,304 sequence reads were generated from 42 samples when targeting the ITS2 and the 18S rDNA, respectively. For sample number 28, only five 18S rDNA reads were generated, and it was thus excluded from further analysis of the 18S rDNA dataset. Sequence trimming and processing resulted in 171,872 ITS2 and 404,281 18S rDNA sequence reads (Suppl. Table S2). Of these, 116,056 ITS2 sequences and 126,051 18S rDNA sequences were of fungal origin. The 18S rDNA sequences generated by co-amplification of the primer pair clustered mainly within the groups of *Telonema*, Alveolata and Stramenopiles.

3.2. Taxonomic composition of the samples

The ITS2 dataset revealed Ascomycota to be the dominant phylum in all sample types, comprising between 72 to 96% of the relative (rel.) sequence abundance, from which 38 – 78% could not be further taxonomically assigned. Dothideomycetes and Eurotiomycetes were the most abundant ascomycete classes under a pH of 8.26 and 7.67, respectively. Basidiomycota were found to be present in all samples but with varying abundance between the two treatments. Up to 22% of the sequences were assigned to Agaricomycetes as the dominant group in samples with a pH of 8.26. In contrast, Microbotryomycetes was the most represented basidiomycete class in samples with a pH of 7.67. No basal fungal lineages were described in the ITS2 dataset, and up to 10% of the sequences could not be assigned beyond the kingdom level (Fig. 1A). All abundant fungal groups in the present-day pH samples with incubation times of up to two weeks were also detected in the positive control (Suppl. Table 3).

In the 18S rDNA dataset, nearly all sequences could be assigned beyond the phylum level, leaving no sequences with an ambiguous taxonomic assignment at higher levels. Ascomycota showed similar

abundance values (74 – 98%) as in the ITS2 dataset, with Eurotiomycetes and Leotiomycetes as dominant groups in both sample types, followed by Dothideomycetes. The percentage of Basidiomycota sequences was relatively small (1 – 7%) and did not exceed more than 2% of the rel. sequence abundance in samples with a pH of 7.67. Using the 18S rDNA dataset, five basal fungal lineages *sensu lato* were detected, namely, Chytridiomycota, Cryptomycota, LKM15, Mucoromycotina and Neocallimastigomycota. In the *in situ* pH treatments, the fraction of sequences assigned to basal fungal lineages declined over time but started with a value of 20% rel. sequence abundance. The positive control was more similar to the present-day pH samples than to the near-future pH samples but contained all abundant groups of both treatments (Suppl. Table 4).

Several fungal groups could not be tested for significant effects of pH and incubation time on their occurrence, as their abundances were too low for reliable results. Thus, only five out of fifteen and eleven out of twenty-seven groups of the ITS2 and 18S rDNA datasets, respectively, were included in this analysis. Of those, the abundance of ten groups was significantly affected by the factor “pH”. With the exceptions of Eurotiomycetes and ambiguous Ascomycota, the remaining eight groups, namely, Cryptomycota, Chytridiomycetes, Microbotryomycetes, Agaricomycetes, Ascomycota *incertae sedis*, Sordariomycetes, Saccharomycetes and Dothideomycetes, had consistently lower abundances under the reduced pH of 7.67 (Table 1).

3.3. ITS-based diversity analyses

Alpha diversity: GLS analysis identified the interaction term of time and pH as significant for Shannon diversity ($F_{3,31}=8.62$, $p=0.0003$). For Simpson diversity, the interaction term was marginally significant ($F_{3,31}=2.66$, $p=0.065$), while time ($F_{3,34}=6.1$, $p=0.0022$) and pH ($F_{1,34}=10.59$, $p=0.0022$) had significant effects. For richness, the best model in terms of residuals included the interaction

term (Table 2). The diversity of the abundant species declined at a pH of 7.67 (Suppl. Table S5, Table 2). In contrast, the richness was higher at a pH of 8.26.

Beta diversity: Bray-Curtis similarity-based PCoA of the ITS2 dataset (Fig. 2) showed a clear separation of samples according to the pH treatment, which is supported by the PERMANOVA analysis (Table 3). However, neither the factor “incubation time” nor the interaction term was found to be significant.

The SIMPER analyses indicated that the OTUs of 15 individual taxa were largely responsible for the observed differences in fungal communities between the samples with different pH levels. The average dissimilarity between the two tested groups was 81.9%. Fourteen of the taxa were part of the core community of the microcosms with an adjusted pH of 7.67. In total, only seven taxa could be identified to the species level, and of those, three were known potential pathogens (Table 4).

3.4. Phylogenetic analysis

Phylogenetic community structure: The unweighted UniFrac-based PCoA did not clearly separate the samples with different pH treatments (Fig. 3A). Similar results were found when using the weighted UniFrac as a metric, but the separation was slightly more pronounced than for the first analysis (Fig. 3B), while incubation time only showed an effect on samples from the first week. This indicates that most of the samples are relatively similar from a phylogenetic perspective and that observed community shifts must be explained based on shifts in taxa abundance, which is in accordance with the non-phylogenetically based PCoA analyses.

Phylogenetic signal analysis: The Phylocom analysis showed significant phylogenetic signals in both community types. The signals always indicated phylogenetic clustering with values of 2.2 for the NRI and 1.2 for the NTI and solely with a value of 1.6 for the NRI in samples with a pH of 8.26 and 7.67, respectively (Fig. 4A). The Nodessig analysis indicated that the phylogenetic signals were

251 mainly caused by the same clades under both pH conditions. The exceptions were significant
252 sequence clustering within clades of Saccharomycetales and Pleosporales exclusively in samples
253 with an adjusted pH (Fig. 4B).

4. Discussion

4.1. Assessing the double-marker gene approach for community analysis

In our approach, we used two marker genes, namely, the ITS and 18S rDNA. The ITS resolves many fungal taxa to the species or genus level but performs less well on basal fungal lineages and lacks phylogenetic power (Schoch et al. 2012). Marine fungal communities are characterized by their (larger) fraction of basal fungi (Panzer et al. 2015, Richards et al. 2015) and many undescribed fungal taxa and groups (Manohar & Raghukumar 2013). Their classification depends on the respective phylogenetic marker applied (Le Calvez et al. 2009, Jones et al. 2011, Nagahama et al. 2011). Indeed, in our study, nearly all fungal sequences were classified to a lower taxonomic level with the 18S rDNA, also revealing the presence of basal fungi. In contrast, the taxonomy of several fungal sequences was only poorly resolved with the ITS, and no basal fungal lineages were detected (Fig. 1).

A phylogenetic-based community approach differs from a non-phylogenetic one with respect to the method used to classify the sequence reads and the metrics used for statistics. The former incorporates the phylogenetic relatedness among taxa into the analyses, which offers further insights into the evolutionary and ecological drivers shaping the community (Horner-Devine & Bohannan 2006, Emerson et al. 2011). In contrast, the standard approach for non-phylogenetic markers, such as the ITS, includes similarity threshold-based sequence clustering (e.g., (Li & Godzik 2006, Edgar 2010)), taxonomic OTU assignment by k-mer similarity search (e.g., (Altschul et al. 1990)) and - most critically - statistics that consider all OTUs as equivalent independently of whether they are phylogenetically highly divergent or not (Faith 1994, Martin 2002). In our approach, we combined the individual advantages of the two marker genes rather than using them in an additive manner as done by Arfi *et al.* (2012) and additionally minimized the primer bias (Hong et al. 2009). Finally, the ITS, possessing good resolution to the species/genus level, was used for the alpha and beta diversity

analyses and for the identification of core species (by SIMPER). In contrast, the 18S rDNA was used to resolve the phylogeny of undescribed taxa/groups and basal fungal lineages to reveal differences in the phylogenetic structure among communities and to test for possible evolutionary conservatism of pH sensitivity in fungal clades.

4.2. Community responses

The mycoplankton community of the microcosms showed a significant response to the pH treatment. However, the communities in the two treatments were phylogenetically highly related, and the observed effect was mainly based on a change in abundance among the dominant taxa rather than a change in the general community composition or extinction of several fungal groups (Fig. 2). Thus, the mycoplankton community can cope with a drop in pH within the tested range, but sensitivity to the pH change differs highly among fungal taxa. The phylogenetic signal analysis indicated no clade-specific sensitivity to one or the other pH (Fig. 4). A phylogenetic signal can occur when, for example, closely related taxa share a character or suite of characters allowing them to be adaptive to the given environmental conditions (Horner-Devine & Bohannan 2006). In the communities in the two pH treatments, small significant clustering effects were observed, but phylogenetic signals were nearly exclusively caused by the same clades in both pH treatments (Fig. 4). This observation points to other factors than the pH causing the clustering effect. In our case, it can be assumed that the habitat-specific conditions of the pelagial at Helgoland Roads are the structuring factors. Panzer *et al.* (2015) recently demonstrated that aquatic fungal community structure is highly influenced by the source habitat, resulting in habitat-specific phylogenetic signals. Thus, the sensitivity to a pH shift observed in our study must be caused by different and (clade-) independent characters; it has, for example, been shown that diverse transcriptional regulatory systems control the response to the environmental pH (Penalva & Arst 2004, Selvig & Alspaugh 2011). In addition, fungi seem to react quite differently to the environmental pH depending on their life stage; for example, the fungal

germination time and outgrowth can be affected by the environmental pH (Magan & Lacey 1984, Porter et al. 1987). For pathogenic fungi with host-independent life stages (e.g., some Cryptomycota or Chytridiomycetes species), responses to pH may depend on the life stage.

4.3. Relevance for food webs

Microbial communities highly influence resource use, disease dynamics and the stability of ecosystem processes (Duffy & Stachowicz 2006, Stachowicz & Byrnes 2006). For example, a pH-dependent loss of fungal diversity in freshwater systems (Wood-Eggenschwiler & Bärlocher 1983, Tolkkinen et al. 2015) can lead to a drastic reduction in fungal leaf breakdown (Baudoin et al. 2008). So far, comparable data for marine fungi are lacking, but recent studies on marine mycoplankton have elucidated their importance in the breakdown of organic matter (Gutierrez et al. 2011) and the control of phytoplankton populations (Gleason et al. 2008, Gutierrez et al. 2016). Thus, zoosporic fungi are part of the pelagic food web, forming a “mycoloop”; herein, they (i) act as a food source for zooplankton grazers, (ii) decompose particulate organic matter, (iii) convert inorganic compounds into organic compounds, and (iv) parasite/hyperparasite on diverse organisms (Kagami et al. 2007, Kagami et al. 2014). In our study, players in the “mycoloop”, namely, Cryptomycota and Chytridiomycetes, showed significant reductions under a reduced pH (Table 1). According to dynamical model predictions, a reduction in zoosporic fungi in pelagic systems would affect the overall food web structure: fewer fungal parasites lead to a higher abundance of large phytoplankton species (>40 µm), which in turn lowers the numbers of smaller phytoplankton species due to resource competition. As zooplankton species only graze on smaller phytoplankton, a reduction in zoosporic fungi has an indirect and negative effect on zooplankton grazers (Miki et al. 2011). The involvement of filamentous fungi in the pelagic food web has so far not been modeled due to missing data. In our study, two of the most prominent groups, Eurotiomycetes and Dothideomycetes, both showed a significant pH effect (Table 1). Gutierrez *et al.* (2011) suggest an important role in the degradation of

326 phytoplankton bloom-derived organic matter for these fungi. Thus, a pH-driven increase in
327 filamentous fungal biomass might have a negative impact on organisms competing for the same food
328 source.

329 Three of the seven key taxa (described to the genus level) that significantly benefited from the pH
330 reduction are pathogenic fungi that cause disease in marine algae (Almeida et al. 2010) and animals
331 (Wirth & Goldani 2012) (Table 4). Pathogenic fungal activities in marine environments may lead to
332 the breakdown of ecosystems (Raghukumar & Ravindran 2012) or cause significant economic losses
333 when commercially important organisms are infected (Gachon et al. 2010, Hatai 2012). Given that
334 pathogenicity is regulated by several factors (e.g., host physiology, species interactions and
335 environmental factors (Fuhrman 2009, Selvig & Alspaugh 2011)), more detailed studies are urgently
336 needed to examine the effects of ocean acidification on this important group.

337 **4.4. Conclusion**

338 Although the marine fungal communities in the two pH treatments were phylogenetically closely
339 related, their taxonomic composition and diversity differed significantly. The observed changes were
340 mainly caused by a restructuring of the abundant species. Sensitivity to the pH reduction was not
341 phylogenetically conserved. The reported pH sensitivity contrasts with the wide pH tolerance
342 observed in marine fungi under controlled conditions. We have only just begun to understand the
343 ecological role of marine fungi, but given the newly discovered role mycoplankton play in organic
344 matter breakdown and phytoplankton bloom dynamics, further studies are urgently needed to place
345 the observed pH-dependent community shift and possible consequences for the marine ecosystem
346 into context.

347 **5. Acknowledgments**

348 We are grateful to Stefanie Thomföhrde for bioinformatic assistance. The authors thank Antje Labes
349 and Lothar Reich for the critical reading of the manuscript. Financial support was provided by the
350 Federal Ministry of Education and Research (BMBF, <https://www.bmbf.de/en/index.html>, FKZ
351 03F0608B) through the project ‘Biological Impacts of Ocean ACIDification’ (BIOACID).

352 **6. Author contributions**

353 Experimental design: EK, AW, GG. Sampling and setup of the experiment: EK. Data analyses: MR,
354 KP. Statistical analyses: GG, LG. Writing of the paper: MR.

355 7. References

- 356 Almeida C, Eguereva E, Kehraus S, Siering C, Konig GM (2010) Hydroxylated sclerosporin
357 derivatives from the marine-derived fungus *Cadophora malorum*. J Nat Prod 73:476-478
- 358 Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J
359 Mol Biol 215:403-410
- 360 Arfi Y, Buee M, Marchand C, Levasseur A, Record E (2012) Multiple markers pyrosequencing
361 reveals highly diverse and host-specific fungal communities on the mangrove trees *Avicennia*
362 *marina* and *Rhizophora stylosa*. FEMS Microbiol Ecol 79:433-444
- 363 Baudoin JM, Guerold F, Felten V, Chauvet E, Wagner P, Rousselle P (2008) Elevated aluminium
364 concentration in acidified headwater streams lowers aquatic hyphomycete diversity and
365 impairs leaf-litter breakdown. Microb Ecol 56:260-269
- 366 Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De Wit P, Sanchez-
367 Garcia M, Ebersberger I, de Sousa F, Amend AS, Jumpponen A, Unterseher M, Kristiansson
368 E, Abarenkov K, Bertrand YJK, Sanli K, Eriksson KM, Vik U, Veldre V, Nilsson RH (2013)
369 Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences
370 of fungi and other eukaryotes for analysis of environmental sequencing data. Methods Ecol
371 Evol 4:914-919
- 372 Blackford JC, Gilbert FJ (2007) pH variability and CO₂ induced acidification in the North Sea. J
373 Marine Syst 64:229-241
- 374 Borneman J, Hartin RJ (2000) PCR primers that amplify fungal rRNA genes from environmental
375 samples. AEM 66:4356-4360
- 376 Bowman BJ, Bowman EJ (1986) H⁺-ATPases from mitochondria, plasma membranes, and vacuoles
377 of fungal cells. J Membrane Biol 94:83-97
- 378 Caldeira K, Wickett ME (2003) Oceanography: Anthropogenic carbon and ocean pH. Nature
379 425:365-365
- 380 Caldeira K, Wickett ME (2005) Ocean model predictions of chemistry changes from carbon dioxide
381 emissions to the atmosphere and ocean. J Geophys Res-Oceans 110:C09S04
- 382 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena
383 AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,
384 Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Tumbaugh PJ,
385 Walters WA, Widmann J, Yatsunenkov T, Zaneveld J, Knight R (2010) QIIME allows analysis
386 of high-throughput community sequencing data. Nat Methods 7:335-336
- 387 Courty PE, Pritsch K, Schlöter M, Hartmann A, Garbaye J (2005) Activity profiling of
388 ectomycorrhiza communities in two forest soils using multiple enzymatic tests. New Phytol
389 167:309-319

390 Dannemiller KC, Reeves D, Bibby K, Yamamoto N, Peccia J (2014) Fungal high-throughput
391 taxonomic identification tool for use with Next-Generation Sequencing (FHiTINGS). J Basic
392 Microb 54:315-321

393 Duffy JE, Stachowicz JJ (2006) Why biodiversity is important to oceanography: potential roles of
394 genetic, species, and trophic diversity in pelagic ecosystem processes. Mar Ecol Prog Ser
395 311:179-189

396 Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. Bioinformatics
397 26:2460-2461

398 Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and
399 speed of chimera detection. Bioinformatics 27:2194-2200

400 Emerson BC, Cicconardi F, Fanciulli PP, Shaw PJA (2011) Phylogeny, phylogeography,
401 phylobetadiversity and the molecular analysis of biological communities. Philosophical
402 Transactions of the Royal Society B-Biological Sciences 366:2391-2402

403 Faith DP (1994) Phylogenetic Pattern and the Quantification of Organismal Biodiversity. Philos T
404 Roy Soc B 345:45-58

405 Fuhrman JA (2009) Microbial community structure and its functional implications. Nature 459:193-
406 199

407 Gachon CMM, Sime-Ngando T, Strittmatter M, Chambouvet A, Kim GH (2010) Algal diseases:
408 spotlight on a black box. Trends Plant Sci 15:633-640

409 Galecki A, Burzykowski T (2013) Linear mixed-effects models using R, Vol. Springer, Berlin
410 Heidelberg

411 Giardine B, Riemer C, Hardison RC, Burhans R, Elnitski L, Shah P, Zhang Y, Blankenberg D, Albert
412 I, Taylor J, Miller W, Kent WJ, Nekrutenko A (2005) Galaxy: A platform for interactive
413 large-scale genome analysis. Genome Res 15:1451-1455

414 Gleason FH, Kagami M, Lefèvre E, Sime-Ngando T (2008) The ecology of chytrids in aquatic
415 ecosystems: roles in food web dynamics. Fungal Biol Rev 22:17-25

416 Goecks J, Nekrutenko A, Taylor J, Team G (2010) Galaxy: a comprehensive approach for supporting
417 accessible, reproducible, and transparent computational research in the life sciences. Genome
418 Biol 11

419 Gutierrez MH, Jara AM, Pantoja S (2016) Fungal parasites infect marine diatoms in the upwelling
420 ecosystem of the Humboldt Current System of central Chile. Environ Microbiol:In press

421 Gutierrez MH, Pantoja S, Tejos E, Quinones RA (2011) The role of fungi in processing marine
422 organic matter in the upwelling ecosystem off Chile. Mar Biol 158:205-219

423 Harvell CD, Mitchell CE, Ward JR, Altizer S, Dobson AP, Ostfeld RS, Samuel MD (2002) Ecology -
424 Climate warming and disease risks for terrestrial and marine biota. Science 296:2158-2162

425 Hatai K (2012) Diseases of fish and shellfish caused by marine fungi. In: Raghukumar C (ed)
426 Biology of marine fungi. Springer, Heidelberg

427 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale
428 PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N,
429 Bradbury RH, Dubi A, Hatziolos ME (2007) Coral reefs under rapid climate change and
430 ocean acidification. *Science* 318:1737-1742

431 Hong SH, Bunge J, Leslin C, Jeon S, Epstein SS (2009) Polymerase chain reaction primers miss half
432 of rRNA microbial diversity. *ISME J* 3:1365-1373

433 Horner-Devine MC, Bohannan BJM (2006) Phylogenetic clustering and overdispersion in bacterial
434 communities. *Ecology* 87:S100-S108

435 Ihrmark K, Bodeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y, Stenlid J,
436 Brandstrom-Durling M, Clemmensen KE, Lindahl BD (2012) New primers to amplify the
437 fungal ITS2 region - evaluation by 454-sequencing of artificial and natural communities.
438 *FEMS Microbiol Ecol* 82:666-677

439 IPCC (2014) Climate change 2014: Mitigation of Climate Change. In: Edenhofer O, Pichs-Madruga
440 R, Sokona Y, Farahani E, Kadner S, Seyboth K, Adler A, Baum I, Brunner S, Eickemeier P,
441 Kriemann B, Savolainen J, Schlömer S, von Stechow C, Zwickel T, Minx JC (eds)
442 Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental
443 Panel on Climate Change. Cambridge University Press, Cambridge, UK

444 Jones MDM, Forn I, Gadelha C, Egan MJ, Bass D, Massana R, Richards TA (2011) Discovery of
445 novel intermediate forms redefines the fungal tree of life. *Nature* 474:200-U234

446 Kagami M, de Bruin A, Ibelings BW, Van Donk E (2007) Parasitic chytrids: their effects on
447 phytoplankton communities and food-web dynamics. *Hydrobiologia* 578:113-129

448 Kagami M, Miki T, Takimoto G (2014) Mycoloop: chytrids in aquatic food webs. *Front Microbiol* 5

449 Krause E, Wichels A, Gimenez L, Gerdt G (2013) Marine fungi may benefit from ocean
450 acidification. *Aquat Microb Ecol* 69:59-67

451 Krause E, Wichels A, Gimenez L, Lunau M, Schilhabel MB, Gerdt G (2012) Small Changes in pH
452 have direct effects on marine bacterial community composition: A microcosm approach.
453 *PLoS ONE* 7:e47035

454 Le Calvez T, Burgaud G, Mahe S, Barbier G, Vandenkoornhuyse P (2009) Fungal diversity in Deep-
455 Sea hydrothermal ecosystems. *AEM* 75:6415-6421

456 Li WZ, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or
457 nucleotide sequences. *Bioinformatics* 22:1658-1659

458 Magan N, Lacey J (1984) Effect of temperature and pH on water relations of field and storage fungi.
459 *T Brit Mycol Soc* 82:71-81

460 Manohar CS, Raghukumar C (2013) Fungal diversity from various marine habitats deduced through
461 culture-independent studies. *FEMS Microbiol Lett* 341:69-78

462 Martin AP (2002) Phylogenetic approaches for describing and comparing the diversity of microbial
463 communities. *AEM* 68:3673-3682

464 Miki T, Takimoto G, Kagami M (2011) Roles of parasitic fungi in aquatic food webs: a theoretical
465 approach. *Freshwater Biol* 56:1173-1183

466 Nagahama T, Takahashi E, Nagano Y, Abdel-Wahab MA, Miyazaki M (2011) Molecular evidence
467 that deep-branching fungi are major fungal components in deep-sea methane cold-seep
468 sediments. *Environ Microbiol* 13:2359-2370

469 Nilsson RH, Wurzbacher C, Bahram M, Coimbra VRM, Larsson E, Tedersoo L, Eriksson J, Ritter
470 CD, Svantesson S, Sanchez-Garcia M, Ryberg M, Kristiansson E, Abarenkov K (2016) Top
471 50 most wanted fungi. *Myckeys*:29-40

472 Orsi W, Biddle JF, Edgcomb V (2013) Deep sequencing of subseafloor eukaryotic rRNA reveals
473 active fungi across marine subsurface provinces. *PLoS ONE* 8

474 Orsi WD, Richards TA, Santoro AE (2015) Cellular maintenance processes that potentially underpin
475 the survival of subseafloor fungi over geological timescales. *Estuar Coast Shelf S* 164:A1-A9

476 Panzer K, Yilmaz P, Weiß M, Reich L, Richter M, Wiese J, Schmaljohann R, Labes A, Imhoff JF,
477 Glöckner FO, Reich M (2015) Identification of habitat-specific biomes of aquatic fungal
478 communities using a comprehensive nearly full-length 18S rRNA dataset enriched with
479 contextual data. *PLoS ONE* 10:e0134377

480 Penalva MA, Arst HN (2004) Recent advances in the characterization of ambient pH regulation of
481 gene expression in filamentous fungi and yeasts. *Annu Rev Microbiol* 58:425-451

482 Pinheiro J, Bates D, DebRoy S, Team RC (2015) *nlme*: Linear and nonlinear mixed effects models.
483 In: R package version 3.1-128, <http://CRAN.R-project.org/package=nlme>.

484 Porter WM, Robson AD, Abbott LK (1987) Field survey of the distribution of vesicular arbuscular
485 mycorrhizal fungi in relation to soil-pH. *J Appl Ecol* 24:659-662

486 Pritsch K, Raidl S, Marksteiner E, Blaschke H, Agerer R, Schloter M, Hartmann A (2004) A rapid
487 and highly sensitive method for measuring enzyme activities in single mycorrhizal tips using
488 4-methylumbelliferone-labelled fluorogenic substrates in a microplate system. *J Microbiol*
489 *Methods* 58:233-241

490 R Core Team (2015) R: A language and environment for statistical computing. In: R Foundation for
491 statistical computing, Vienna, Austria

492 Raghukumar C, Ravindran J (2012) Fungi and their role in corals and coral reef ecosystems. In:
493 Raghukumar C (ed) *Biology of marine fungi*. Springer, Heidelberg

494 Rama T, Davey M, Norden J, Halvorsen R, Błaalid R, Mathiassen G, Alsos I, Kauserud H (2016)
 495 Fungi sailing the Arctic Ocean: Speciose communities in North Atlantic driftwood as
 496 revealed by high-throughput amplicon sequencing. *Microb Ecol* 72:295-304

497 Richards TA, Leonard G, Mahé F, del Campo J, Romac S, Jones MDM, Maguire F, Dunthorn M, De
 498 Vargas C, Massana R, Chambouvet A (2015) Molecular diversity and distribution of marine
 499 fungi across 130 European environmental samples. *Proc R Soc Lond [Biol]* 282

500 Sabine CL, Feely RA, Gruber N, Key RM, Lee K, Bullister JL, Wanninkhof R, Wong CS, Wallace
 501 DWR, Tilbrook B, Millero FJ, Peng TH, Kozyr A, Ono T, Rios AF (2004) The oceanic sink
 502 for anthropogenic CO₂. *Science* 305:367-371

503 Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Bolchacova E,
 504 Voigt K, Crous PW, Miller AN, Wingfield MJ, Aime MC, An KD, Bai FY, Barreto RW,
 505 Begerow D, Bergeron MJ, Blackwell M, Boekhout T, Bogale M, Boonyuen N, Burgaz AR,
 506 Buyck B, Cai L, Cai Q, Cardinali G, Chaverri P, Coppins BJ, Crespo A, Cubas P, Cummings
 507 C, Damm U, de Beer ZW, de Hoog GS, Del-Prado R, B D, Dieguez-Urbeondo J, Divakar
 508 PK, Douglas B, Duenas M, Duong TA, Eberhardt U, Edwards JE, Elshahed MS, Fliegerova
 509 K, Furtado M, Garcia MA, Ge ZW, Griffiths GW, Griffiths K, Groenewald JZ, Groenewald
 510 M, Grube M, Gryzenhout M, Guo LD, Hagen F, Hambleton S, Hamelin RC, Hansen K,
 511 Harrold P, Heller G, Herrera G, Hirayama K, Hirooka Y, Ho HM, Hoffmann K, Hofstetter V,
 512 Hognabba F, Hollingsworth PM, Hong SB, Hosaka K, Houbraken J, Hughes K, Huhtinen S,
 513 Hyde KD, James T, Johnson EM, Johnson JE, Johnston PR, Jones EB, Kelly LJ, Kirk PM,
 514 Knapp DG, Koljalg U, GM K, Kurtzman CP, Landvik S, Leavitt SD, Liggenstoffer AS,
 515 Liimatainen K, Lombard L, Luangsa-Ard JJ, Lumbsch HT, Maganti H, Maharachchikumbura
 516 SS, Martin MP, May TW, McTaggart AR, Methven AS, Meyer W, Moncalvo JM,
 517 Mongkolsamrit S, Nagy LG, Nilsson RH, Niskanen T, Nyilasi I, Okada G, Okane I, Olariaga
 518 I, Otte J, Papp T, Park D, Petkovits T, Pino-Bodas R, Quaedvlieg W, Raja HA, Redecker D, T
 519 R, Ruibal C, Sarmiento-Ramirez JM, Schmitt I, Schussler A, Shearer C, Sotome K, Stefani
 520 FO, Stenroos S, Stielow B, Stockinger H, Suetrong S, Suh SO, Sung GH, Suzuki M, Tanaka
 521 K, Tedersoo L, Telleria MT, Tretter E, Untereiner WA, Urbina H, Vagvolgyi C, Vialle A, Vu
 522 TD, Walther G, Wang QM, Wang Y, Weir BS, Weiss M, White MM, Xu J, Yahr R, Yang
 523 ZL, Yurkov A, Zamora JC, Zhang N, Zhuang WY, Schindel D, Consortium FB (2012)
 524 Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode
 525 marker for Fungi. *Proc Natl Acad Sci U S A* 109:6241-6246

526 Selvig K, Alspaugh JA (2011) pH response pathways in fungi: Adapting to host-derived and
 527 environmental signals. *Mycobiology* 39:249-256

528 Stachowicz JJ, Byrnes JE (2006) Species diversity, invasion success, and ecosystem functioning:
 529 Disentangling the influence of resource competition, facilitation, and extrinsic factors. *Mar*
 530 *Ecol Prog Ser* 311:251-262

531 Taylor JD, Cunliffe M (2016) Multi-year assessment of coastal planktonic fungi reveals
 532 environmental drivers of diversity and abundance. *ISME J* 10:2118-2128

533 Tolkkinen M, Mykra H, Annala M, Markkola AM, Vuori KM, Muotka T (2015) Multi-stressor
 534 impacts on fungal diversity and ecosystem functions in streams: natural vs. anthropogenic
 535 stress. *Ecology* 96:672-683

536 Venables WN, Ripley BD (2002) Modern applied statistics with S., Vol. Springer-Verlag, New York

537 Verdugo P, Alldredge AL, Azam F, Kirchman DL, Passow U, Santschi PH (2004) The oceanic gel
538 phase: a bridge in the DOM-POM continuum. Mar Chem 92:67-85

539 Webb CO, Ackerly DD, Kembel SW (2008) Phylocom: software for the analysis of phylogenetic
540 community structure and trait evolution. Bioinformatics 24:2098-2100

541 Webster NS, Negri AP, Flores F, Humphrey C, Soo R, Botte ES, Vogel N, Uthicke S (2013) Near-
542 future ocean acidification causes differences in microbial associations within diverse coral
543 reef taxa. Env Microbiol Rep 5:243-251

544 White TJ, Bruns T, Lee S, Taylor JW (1990) Amplification and direct sequencing of fungal
545 ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ
546 (eds) PCR Protocols: A guide to methods and applications. Academic Press, Inc., New York,
547 USA

548 Wiltshire KH, Malzahn AM, Wirtz K, Greve W, Janisch S, Mangelsdorf P, Manly BFJ, Boersma M
549 (2008) Resilience of North Sea phytoplankton spring bloom dynamics: An analysis of long-
550 term data at Helgoland Roads. Limnol Oceanogr 53:1294-1302

551 Wirth F, Goldani LZ (2012) Epidemiology of *Rhodotorula*: An emerging pathogen. Interdiscip
552 Perspect Infect Dis:465717

553 Witt V, Wild C, Anthony KRN, Diaz-Pulido G, Uthicke S (2011) Effects of ocean acidification on
554 microbial community composition of, and oxygen fluxes through, biofilms from the Great
555 Barrier Reef. Environ Microbiol 13:2976-2989

556 Wood-Eggenschwiler S, Bärlocher F (1983) Aquatic hyphomycetes in sixteen streams in France,
557 Germany and Switzerland. T Brit Mycol Soc 81:371-379

558 Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and extensions in
559 ecology with R, Vol. Springer, Heidelberg

560

561

562 **Table 1: Generalized modeling framework test of the effects of pH and week on the 13 most abundant fungal groups.** For several
563 groups, the test was not applicable due to low abundance (n/a). Displayed are the average abundances of the tested fungal groups at a pH of
564 8.26 and 7.67 as observed in the two different datasets, ITS and 18S rDNA. The p-value indicates whether the pH effect had a significant
565 effect on the occurrence of the fungal group and whether this was dependent on incubation time. Finally, the trend in abundance within the
566 two pH treatments is given. Av. Abund., average abundance; SD, standard deviation; P, p-value; L, pH 7.67; H, pH 8.26; ns, not significant.

Fungal group	Av. Abund. \pm (SD) at pH 8.26 (ITS)	Av. Abund. \pm (SD) at pH 7.67 (ITS)	p (ITS)	Av. Abund. \pm (SD) at pH 8.26 (18S rDNA)	Av. Abund. \pm (SD) at pH 7.67 (18S rDNA)	P (18S rDNA)	Trend in abund. over the two pH treatments
Cryptomycota	0	0	n/a	350 \pm 25.1	23 \pm 1.1	4.3x10 ⁻¹⁴	L<H
Chytridiomycetes	0	0	n/a	184 \pm 5.1	61 \pm 2.8	1.47x10 ⁻⁰⁸	L<H
Microbotryomycetes	16 \pm 1.3	58 \pm 2.5	10 ⁻⁴	44 \pm 2.4	35 \pm 1.4	8.8x10 ⁻³ (for week 4) 4.2x10 ⁻² (for week 1)	L>H L<H
Agaricomycetes	92 \pm 3.6	1 \pm 0	10 ⁻⁸	66 \pm 5.7	14 \pm 1.3	8x10 ⁻⁴	L<H
Ambiguous Ascomycota	371 \pm 8.1	582 \pm 4.9	2x10 ⁻⁴	16 \pm 1.2	0	n/a	L>H
Ascomycota <i>Incertae Sedis</i>	2 \pm 0	0	n/a	42 \pm 2.1	16 \pm 1	2x10 ⁻³	L<H
Sordariomycetes	15 \pm 1.2	4 \pm 0	n/a	146 \pm 14	80 \pm 1.9	0.01 (for week 3)	L<H
Saccharomycetes	29 \pm 3.9	0	n/a	62 \pm 6.2	4 \pm 0	3.57x10 ⁻⁰⁵	L<H
Leotiomycetes	42 \pm 5.9	2 \pm 0	n/a	2163 \pm 13.3	2174 \pm 19.4	0.95	ns
Lecanoromycetes	5 \pm 0.5	0	n/a	33 \pm 1.2	38 \pm 1.3	0.41	ns
Dothideomycetes	133 \pm 10.2	7 \pm 0.8	n/a	360 \pm 14.8	201 \pm 3.5	0.0	L<H
Eurotiomycetes	1 \pm 0	133 \pm 2.7	n/a	1520 \pm 26.5	201 \pm 3.5	2.25x10 ⁻⁰⁵	L>H

567

568 **Table 2: Results of generalized linear model analysis testing the effects of incubation time and pH on the richness.** Test included the
569 examination of negative binomial and Poisson error structure. Likelihood ratio tests (LR) and associated (Chi squared-based) p-values
570 compare the top two models, testing for the significance of the interaction term. The LR uses the degrees of freedom of the model with the
571 interaction (df=31) and the one without the interaction (df=34) to calculate the difference (df=3) and then the LR(3)=15.46. AIC: Akaike
572 information criterion; LR: likelihood ratio test; p: p-value (Chi squared-based).

Fixed structure	Error structure	AIC	LR	p
Time + pH + TxpH	Negative binomial	341	15.46	0.0014
Time + pH	Negative binomial	351		
Time + pH + TxpH	Poisson	401		
Time + pH	Poisson	467		

573

574 **Table 3: PERMANOVA main test of fungal community composition based on ITS2-based Bray-Curtis dissimilarities of operational**
575 **taxonomic units.** Displayed are tests for the factors “pH” and “incubation time”, their interaction and the partitioning of multivariate
576 variation. p-values were obtained using type III sums of squares and 9999 permutations under the full model. *, statistical significance (p <
577 0.05); d.f.: degrees of freedom; SS: sums of squares; Sq. root: square root of the component of variation attributable to that factor in the
578 model in units of Bray-Curtis dissimilarity.

Sources of variation	d.f.	SS	Pseudo-F	Sq.root
pH*	1	28308	15,177	36,925
Time	3	8301	14,835	96,348
pH × time	3	7791	13,924	12,275
Residuals	31	57821	43,188	
Total	38	1025200000		

579

580 **Table 4: SIMPER analysis of the ITS2 dataset.** Operational taxonomic units (OTUs) contributing most to the observed community
581 dissimilarity among the two pH sample types. Only OTUs with more than 1% contribution to the observed community dissimilarities are
582 listed. Av. Abund., average abundance; Av. Diss, average dissimilarity.

Species	Av. Abund. at pH 8.26	Av. Abund. at pH 7.67	Av. Diss.	COD (%)	Phylum	Subphylum	Class	Order
<i>Cyphellophora reptans</i>	0.49	32.78	13.39	16.33	Ascomycota	Pezizomycotina	Eurotiomycetes	Chaetothyriales
<i>Pezizomycotina</i> sp.	7.85	22.94	6.6	8.06	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Sporobolomyces symmetricus</i>	0.38	7.56	2.85	3.48	Basidiomycota	Pucciniomycotina	Cystobasidiomycetes	Sporidiobolales
<i>Cadophora malorum</i>	1.16	6.87	2.84	3.46	Ascomycota	Pezizomycotina	Leotiomycetes	Helotiales
<i>Sakaguchia dacryoidea</i>	1.04	4.87	2.14	2.6	Basidiomycota	Pucciniomycotina	Cystobasidiomycetes	Incertae sedis
<i>Pezizomycotina</i> sp.	2.13	6.27	1.82	2.22	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Eurotiomycetes</i> sp.	0.11	3.63	1.47	1.8	Ascomycota	Pezizomycotina	Eurotiomycetes	Incertae sedis
<i>Pezizomycotina</i> sp.	1.37	4.35	1.29	1.58	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Neophaeosphaeria quadrisepata</i>	2.28	1.2	1.26	1.54	Ascomycota	Pezizomycotina	Dothideomycetes	Pleosporales
<i>Pezizomycotina</i> sp.	1.15	3.95	1.25	1.53	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Pezizomycotina</i> sp.	1.22	3.89	1.21	1.48	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Pezizomycotina</i> sp.	0.77	2.82	0.91	1.11	Ascomycota	Pezizomycotina	Incertae sedis	Incertae sedis
<i>Eurotiomycetes</i> sp.	0	2.2	0.9	1.1	Ascomycota	Pezizomycotina	Eurotiomycetes	Incertae sedis
<i>Acremonium butyri</i>	0.76	2.57	0.88	1.07	Ascomycota	Pezizomycotina	Sordariomycetes	Hypocreales

<i>Rhodotorula aurantiaca</i>	0.21	2.3	0.85	1.04	<i>Basidiomycota</i>	<i>Pucciniomycotina</i>	<i>Cystobasidiomycetes</i>	Incertae sedis
-------------------------------	------	-----	------	------	----------------------	-------------------------	----------------------------	----------------

583

584 **Figure legends**


585 **Figure 1: Taxonomic assemblage of fungal communities in the two tested pH sample types over**
586 **four weeks of incubation.** Taxonomic composition based on the (A) ITS and (B) 18S rDNA dataset.
587 Fungal groups in bold showed a significant pH response: *, reduced abundance at a pH of 7.67; \$,
588 increased abundance at a pH of 7.67; #, not all time points showed consistent effects. For more detail,
589 see Table 2. 1-4, weeks of incubation. Bottom to dashed lines: *Ascomycota*; dashed to dotted lines:
590 *Basidiomycota*; dashed lines to top: basal fungal lineages *sensu lato*.

591 **Figure 2: Principal coordinate analysis (PCoA) of the ITS2 dataset.** The test is based on square
592 root-transformed data using Bray-Curtis dissimilarities of operational taxonomic unit counts.

593 **Figure 3: Ordination analysis of phylogenetic distances revealing phylogenetic relatedness**
594 **among communities.** PCoAs are based on (A) unweighted and (B) weighted UniFrac metrics using
595 the 18S rDNA data.

596 **Figure 4: Identification of phylogenetic signals via Phylocom analysis testing for clade-specific**
597 **adaptation to a pH of 8.26 and 7.67.** (A) Comstruct analysis revealed significant phylogenetic
598 clustering in samples for both pH treatments. (B) Nodesig analysis identified the fungal groups that
599 caused the observed phylogenetic signals. The schematic phylogenetic tree depicts only the fungal
600 clades that significantly contributed to the observed phylogenetic clustering. Identified significant
601 clustering was assigned to class/subphylum (left) and order levels (right). NRI, net relatedness index;
602 NTI, nearest taxon index; blue, clustering effect present in both pH sample types; red, clustering
603 effect observed only in the sample type with a pH of 8.26; green, effect only observed in sample type
604 with a pH of 7.67.


605 **Figure 1:**



606

607 **Figure 2:**


608



609


610 **Figure 3:**

611



612

A)



B)

